arXiv:hep-ph/9908346v5 3 Jun 2000

Polarized $\Lambda_b \to \Lambda_c l\bar{\nu}$ decay in perturbative QCD

Hsien-Hung Shih* and Shih-Chang Lee†
Institute of Physics, Academia Sinica, Taipei, Taiwan 105, Republic of China
Hsiang-nan Li‡
Department of Physics, National Cheng-Kung University, Tainan, Taiwan 701, Republic of China
Theory Group, KEK, Tsukuba, Ibaraki 305, Japan

(February 1, 2008)

We study the polarized Λ_b baryon semileptonic decay $\Lambda_b \to \Lambda_c l\bar{\nu}$ using perturbative QCD factorization theorem. Employing the heavy baryon transition form factors calculated in our previous work, we predict the asymmetry parameter $\alpha(\hat{e}) \equiv (N_+ - N_-)/(N_+ + N_-) = -0.31\mathbf{P}\cdot\hat{e}$, where $\mathbf{P}$ is the Λ_b baryon polarization, $\hat{e}$ a unit vector perpendicular to the Λ_b baryon momentum, N_+ (N_-) the number of leptons whose momenta possess positive (negative) components along $\hat{e}$. This result is useful for extracting the transverse polarization of a Λ_b baryon from data of exclusive semileptonic decays.

I. INTRODUCTION

Recently, we have developed perturbative QCD (PQCD) factorization theorem for exclusive heavy baryon decays [1,2]. In this approach transition form factors are expressed as convolutions of hard heavy quark decay amplitudes with heavy baryon wave functions. The former are calculable in perturbation theory in the kinematic region where large momentum transfer is involved. The latter, absorbing nonperturbative dynamics of decay processes, must be obtained by means outside the PQCD regime. Large logarithmic corrections are organized by Sudakov resummation and renormalization-group equations to improve perturbative expansions. Since wave functions are universal, they can be determined once for all, and then employed to make predictions for other modes containing the same heavy baryons. With this prescription for nonperturbative wave functions, PQCD factorization theorem possesses predictive power.

In this paper we shall apply the above PQCD formalism to polarized Λ_b baryon decays. It is well known that hyperons produced in hadronic machines through strong interaction are usually polarized. However, the mechanism responsible for such a polarization is still unclear. Hence, it is interesting to investigate if similar phenomena happen for heavy baryons produced in proton-proton and proton-anti-proton collisions at various energies. Since b-baryon events in hadronic productions are often selected by the presence of high energy leptons, we shall focus on studying the Λ_b baryon polarization through the semileptonic decays. The longitudinal polarization of a Λ_b baryon produced from Z^0 decays [3,4] could be determined by measuring the ratio of the average electron (muon) energy to the average neutrino energy in the inclusive semileptonic decays [5,6]. This ratio, obtained in the laboratory frame, is then related to the ratio of the corresponding energies in the rest frame of the Λ_b baryon.

When heavy baryons are produced via strong interaction, they could possess transverse polarization perpendicular to baryon momenta. In the present work we shall concentrate on the determination of the transverse polarization of a Λ_b baryon from the angular distribution of leptons produced in the $\Lambda_b \to \Lambda_c l\bar{\nu}$ decay. For semileptonic decays, it is usually not possible to reconstruct fully the momentum of the parent baryon. In order to determine the heavy baryon polarization through semileptonic events, we propose to measure the asymmetry parameter $\alpha \equiv (N_+ - N_-)/(N_+ + N_-)$, where N_+ (N_-) is the lepton number emitted above and below a plane that contains the Λ_b baryon momentum. This quantity is invariant under a longitudinal boost and can be simply evaluated in the rest frame of the Λ_b baryon. Using the $\Lambda_b \to \Lambda_c$ transition form factors obtained from PQCD factorization theorem [2], we derive the relation between α and the transverse polarization $\mathbf{P}\cdot\hat{e}$ of the Λ_b baryon, $\alpha = -0.31\mathbf{P}\cdot\hat{e}$, where $\hat{e}$ is a unit vector perpendicular to the Λ_b baryon momentum. Comparing this relation with experimental data of α, one may extract the Λ_b baryon transverse polarization.

It has been shown that PQCD analyses for $\Lambda_b \to \Lambda_c$ transition form factors are reliable at the maximal recoil of the Λ_c baryon [2]. The percentages of the full contribution to the form factors, that arise from the short-distance region with $\alpha_s/\pi < 0.2$-0.5, are listed in Table I. It is observed that for the velocity transfer $\rho \sim 1.4$, about 60% of the full contribution comes from the region with $\alpha_s/\pi < 0.3$. Though the results are not completely perturbative, it makes sense to estimate the form factors at the maximal recoil using PQCD, and to give predictions that can be tested by future experiments.

---

*hhshih@phys.sinica.edu.tw

†phsclee@ccvax.sinica.edu.tw

‡hnli@mail.ncku.edu.tw

Extrapolating PQCD results to small ρ under the requirement of heavy quark symmetry (HQS) [7], we have determined the behaviors of the form factors in the whole range of the velocity transfer. This extrapolation leads to the reasonable branching ratio $B(\Lambda_b \to \Lambda_c l\bar{\nu}) = 2 \sim 3\%$, which is consistent with the experimental upper bound of the branching ratio from the data $B(\Lambda_b \to \Lambda_c l\bar{\nu} + X) = (8.27 \pm 3.38)\%$ [8]. The $\Lambda_b \to \Lambda_c$ transition form factors have been evaluated by means of overlap integrals of infinite-momentum-frame wave functions [9,10], relativistic quark models [11], Bethe-Salpeter equations [12] and QCD sum rules [13–15]. Most of the analyses led to the branching ratios about or below 6%. Our result is close to the result $(3.4 \pm 0.6)\%$ derived in [13].

II. ASYMMETRY PARAMETER

We first define the $\Lambda_b \to \Lambda_c$ transition form factors, and refer their explicit factorization formulas to [2]. The amplitude for the semileptonic decay $\Lambda_b \to \Lambda_c l\bar{\nu}$ is written as

$$\mathcal{M} = \frac{G_F}{\sqrt{2}} V_{cb}\, \bar{l}\gamma^\mu(1-\gamma_5)\nu_l\, \langle \Lambda_c(p')|\bar{c}\gamma_\mu(1-\gamma_5)b|\Lambda_b(p)\rangle\ , \tag{1}$$

where G_F is the Fermi coupling constant, V_{cb} the Cabibbo-Kobayashi-Maskawa (CKM) matrix element, p and p' the Λ_b and Λ_c baryon momenta, respectively. QCD dynamics is contained in the hadronic matrix element

$$\begin{aligned}\mathcal{H}_\mu &\equiv \langle \Lambda_c(p')|\bar{c}\gamma_\mu(1-\gamma_5)b|\Lambda_b(p)\rangle\ ,\\ &= \bar{\Lambda}_c(p')[f_1(q^2)\gamma^\mu - if_2(q^2)\sigma^{\mu\nu}q_\nu + f_3(q^2)q^\mu]\Lambda_b(p)\\ &\quad +\bar{\Lambda}_c(p')[g_1(q^2)\gamma^\mu - ig_2(q^2)\sigma^{\mu\nu}q_\nu + g_3(q^2)q^\mu]\gamma_5\Lambda_b(p)\ .\end{aligned} \tag{2}$$

In the second line $\mathcal{H}_\mu$ has been expressed in terms of six form factors f_i and g_i, where $\Lambda_b(p)$ and $\Lambda_c(p')$ are the Λ_b and Λ_c baryon spinors, respectively, and the variable q denotes $q = p - p'$. In the case of massless leptons with $q_\mu \bar{l}\gamma^\mu(1-\gamma_5)\nu_l = 0$, f_3 and g_3 do not contribute. Since the contributions from f_2 and g_2 are small, we shall consider only f_1 and g_1 below.

The polarization density matrix for a Λ_b baryon with polarization P is written as

$$\varrho_{KK'} = \frac{1}{2}P^a(\sigma_a)_{KK'}\ ,\quad \sigma_0 = I\ . \tag{3}$$

For the purpose of normalization, we usually choose $P^0 = 1$. The hadronic matrix element with the helicity of the initial- and final-state baryons specified is expressed as

$$\mathcal{H}_\mu(K,J) = \bar{\Lambda}_c(p',J)[f_1(\rho)\gamma_\mu + g_1(\rho)\gamma_\mu\gamma_5]\Lambda_b(p,K)\ , \tag{4}$$

which leads to the covariant density matrix

$$\mathcal{H}_{\mu\nu} = \sum_{K,K',J} \mathcal{H}_\mu(K,J)\varrho_{KK'}\mathcal{H}^*_\nu(K',J)\ . \tag{5}$$

The differential decay rate is written as

$$d\Gamma = \frac{1}{2M_{\Lambda_b}}|\mathcal{M}_p|^2(2\pi)^4\delta^4(p-p'-p_l-p_\nu)\frac{d^3\mathbf{p}'}{(2\pi)^3 2E'}\frac{d^3\mathbf{p}_l}{(2\pi)^3 2E_l}\frac{d^3\mathbf{p}_\nu}{(2\pi)^3 2E_\nu}\ , \tag{6}$$

where p_l (p_ν) is the lepton (neutrino) momentum. The matrix element square is given by

$$|\mathcal{M}_p|^2 = \frac{G_F^2}{2}|V_{cb}|^2\mathcal{H}_{\mu\nu}\mathcal{L}^{\mu\nu}\ , \tag{7}$$

with the leptonic covariant tensor,

$$\mathcal{L}^{\mu\nu} = 8\left[p_l^\mu p_\nu^\nu + p_l^\nu p_\nu^\mu - p_l\cdot p_\nu g^{\mu\nu} + i\epsilon^{\mu\nu\alpha\beta}p_{l\alpha}p_{\nu\beta}\right]\ . \tag{8}$$

Define the velocity transfer ρ, which is related to q^2 via

$$\rho = \frac{M_{\Lambda_b}^2 + M_{\Lambda_c}^2 - q^2}{2M_{\Lambda_b}M_{\Lambda_c}} \ , \tag{9}$$

with M_{Λ_b} and M_{Λ_c} being the Λ_b and Λ_c baryon masses, respectively. Performing the phase space integrations straightforwardly, Eq. (6) reduces to

$$\frac{d^4\Gamma}{dE' dE_l d\cos\theta_c d\Omega_l} = \frac{1}{256\pi^5 M_{\Lambda_b}} \frac{1}{\sin\theta_c \sin\theta_l \sin\bar{\phi}} |\mathcal{M}_p|^2 \ , \tag{10}$$

where $E' = \rho M_{\Lambda_c}$ (E_l) is the Λ_c baryon (lepton) energy, θ_c (θ_l) the polar angle of the Λ_c baryon momentum $\mathbf{p}'$ (the lepton momentum $\mathbf{p}_l$) against the Λ_b baryon polarization, ϕ_l the azimuthal angle of $\mathbf{p}_l$, and $d\Omega_l = d\cos\theta_l d\phi_l$ the solid angle. The azimuthal angle $\bar{\phi}$ and the angle $\bar{\theta}$ between $\mathbf{p}'$ and $\mathbf{p}_l$ satisfy the relations

$$\cos\bar{\phi} = \frac{1}{\sin\theta_c \sin\theta_l}[\cos\bar{\theta} - \cos\theta_c \cos\theta_l] \ , \tag{11}$$

$$\cos\bar{\theta} = \frac{M_{\Lambda_b}^2 + M_{\Lambda_c}^2 - 2M_{\Lambda_b}E' - 2M_{\Lambda_b}E_l + 2E'E_l}{2\sqrt{E'^2 - M_{\Lambda_c}^2}E_l} \ . \tag{12}$$

The kinematic ranges of E' and E_l are

$$\begin{aligned} M_{\Lambda_c} \leq & E' \leq \frac{M_{\Lambda_b}^2 + M_{\Lambda_c}^2}{2M_{\Lambda_b}} \ , \\ \frac{1}{2}\left[M_{\Lambda_b} - E' - \sqrt{E'^2 - M_{\Lambda_c}^2}\right] \leq & E_l \leq \frac{1}{2}\left[M_{\Lambda_b} - E' + \sqrt{E'^2 - M_{\Lambda_c}^2}\right] \ . \end{aligned} \tag{13}$$

Requiring $\cos^2\bar{\phi} \leq 1$, it is easy to derive the ranges of $\cos\theta_c$ and $\cos\theta_l$ from Eq. (11),

$$\begin{aligned} & |\cos\theta_l| \leq 1 \ , \\ \cos(\theta_l + \bar{\theta}) \leq \ & \cos\theta_c \leq \cos(\theta_l - \bar{\theta}) \ . \end{aligned} \tag{14}$$

The projection operator associated with a spin-1/2 particle of mass m is written as [16]

$$u(p,K)\bar{u}(p,K') = \frac{1}{2}(m + \not{p})[(I)_{K'K} + \gamma_5 \gamma_\mu n_i^\mu (\sigma^i)_{K'K}] \ , \tag{15}$$

where I is the 2×2 unit matrix, K and K' represent the helicities $\pm\frac{1}{2}$, and σ^i are the Pauli matrices. In the rest frame of a Λ_b baryon with $p = (M_{\Lambda_b}, \mathbf{0})$, we have the boost vectors $n_i^j = \delta_i^j$, $i,j = 1,2,3$. Using Eqs. (3), (5), and (15), we obtain the expression of $|\mathcal{M}_p|^2$,

$$|\mathcal{M}_p|^2 = 8M_{\Lambda_b} G_F^2 |V_{cb}|^2 [F_0(E',E_l) + F_1(E',E_l)\mathbf{p}' \cdot \mathbf{P} + F_2(E',E_l)\mathbf{p}_l \cdot \mathbf{P}] \ , \tag{16}$$

with the functions

$$\begin{aligned} F_0 = & -(M_{\Lambda_b} - E' - E_l)[M_{\Lambda_b}^2 + M_{\Lambda_c}^2 - 2(E' + E_l)M_{\Lambda_b}](f_1 - g_1)^2 \\ & + E_l(M_{\Lambda_b}^2 - M_{\Lambda_c}^2 - 2M_{\Lambda_b}E_l)(f_1 + g_1)^2 - M_{\Lambda_c}(M_{\Lambda_b}^2 + M_{\Lambda_c}^2 - 2M_{\Lambda_b}E')(f_1^2 - g_1^2) \ , \\ F_1 = & [M_{\Lambda_b}^2 + M_{\Lambda_c}^2 - 2(E' + E_l)M_{\Lambda_b}](f_1 - g_1)^2 + 2M_{\Lambda_c}E_l(f_1^2 - g_1^2) \ , \\ F_2 = & [M_{\Lambda_b}^2 + M_{\Lambda_c}^2 - 2(E' + E_l)M_{\Lambda_b}](f_1 - g_1)^2 - (M_{\Lambda_b}^2 - M_{\Lambda_c}^2 - 2M_{\Lambda_b}E_l)(f_1 + g_1)^2 \\ & + 2M_{\Lambda_c}(M_{\Lambda_b} - E')(f_1^2 - g_1^2) \ , \end{aligned} \tag{17}$$

and the scalar products

$$\mathbf{p}' \cdot \mathbf{P} = \sqrt{E'^2 - M_{\Lambda_c}^2}\ |\mathbf{P}| \cos\theta_c \ , \quad \mathbf{p}_l \cdot \mathbf{P} = E_l\ |\mathbf{P}| \cos\theta_l \ . \tag{18}$$

The first term F_0 is associated with the unpolarized Λ_b baryon decay. Integrating out $\cos\theta_c$ according to the range in Eq. (14), Eq. (10) becomes

$$\frac{d^3\Gamma}{dE' dE_l d\Omega_l} = \frac{G_F^2}{16\pi^4}|V_{cb}|^2[F_0 + (M_{\Lambda_c}\ \sqrt{\rho^2-1}\cos\bar{\theta}F_1 + E_l F_2)\ \hat{\mathbf{p}}_l \cdot \mathbf{P}]\ , \tag{19}$$

where $\hat{\mathbf{p}}_l = \mathbf{p}_l/|\mathbf{p}_l|$ is the unit vector along the lepton momentum.

To obtain the decay rate, we need the information of the form factors $f_1(\rho)$ and $g_1(\rho)$ in the whole range of ρ. f_1 and g_1 have been evaluated in [2] by adopting the CKM matrix element $V_{cb} = 0.04$, the masses $M_{\Lambda_b} = 5.624$ GeV and $M_{\Lambda_c} = 2.285$ GeV. As shown in Table I, the contribution from the short-distance region with $\alpha_s/\pi < 0.3$ becomes dominant gradually when ρ increases. The PQCD analysis can be regarded as being self-consistent at the maximal $\rho \sim 1.4$, for which the perturbative contribution amounts to about 60% of the full contribution. We then extrapolated the PQCD predictions to the small ρ region under the requirement of HQS [7]. Assuming the parametrization,

$$f_1(\rho) = \frac{c_f}{\rho^{\alpha_f}}\ , \quad g_1(\rho) = \frac{c_g}{\rho^{\alpha_g}}\ , \tag{20}$$

we have obtained the constants $c_f = 1.32$ and $c_g = -1.19$, and the powers $\alpha_f = 5.18$ and $\alpha_g = 5.14$ by fitting Eq. (20) the PQCD results at large ρ [2]. The values of the form factors at zero recoil, $f_1(1) = 1.32$ and $g_1(1) = -1.19$, are close to those derived based on wave function overlap integrals [9]. The approximate equality of f_1 and $|g_1|$ is consistent with the conclusion drawn from heavy quark effective theory (HQET) [17,18].

We emphasize that there is no conflict between PQCD and HQET, which has been applied to heavy hadron decays successfully. If a hadron is indeed very heavy, a small portion of its mass used for the energy release involved in decay processes can guarantee the applicability of PQCD. That is, the decay processes move into the perturbative regime quickly once the velocity transfer is slightly greater than unity. In [2] we have made a quantitative investigation to find out how far from the zero recoil PQCD is applicable to Λ_b baryon decays. Besides, HQET gives the relations among various transition form factors, while PQCD can be employed to calculate the behaviors of these form factors near the high end of the velocity transfer. We have also performed the heavy quark expansion as constructing the heavy baryon wave functions. Hence, the two approaches in fact complement each other.

To extract the transverse polarization of a Λ_b baryon, we propose to measure a lepton asymmetry parameter in the laboratory frame. Choose a unit vector $\hat{e}$ perpendicular to the Λ_b baryon momentum, whose direction is known if the interaction point of hadron beams as well as the decay vertex of the Λ_b baryon are observed. The lepton asymmetry parameter $\alpha(\hat{e})$ with respect to $\hat{e}$ is defined as

$$\alpha(\hat{e}) = \frac{N_+ - N_-}{N_+ + N_-}\ , \tag{21}$$

where $N_+(N_-)$ is the number of leptons with positive (negative) values of $\hat{\mathbf{p}}_l \cdot \hat{e}$. Since $\alpha(\hat{e})$ is invariant under a boost along the Λ_b baryon momentum, we can simply calculate $\alpha(\hat{e})$ in the rest frame of the Λ_b baryon, *i.e.*, in the above framework, and derive its relation to the Λ_b baryon polarization.

Integrating over the variables E' (or ρ) and E_l in Eq. (19), we arrive at

$$\frac{d\Gamma}{d\Omega_l} = \frac{M_{\Lambda_c}G_F^2}{16\pi^4}|V_{cb}|^2\ (G_0 + G_1\ \hat{\mathbf{p}}_l \cdot \mathbf{P})\ , \tag{22}$$

with the integrals

$$\begin{aligned} G_0 &= \int d\rho dE_l\ F_0(E', E_l)\ , \\ G_1 &= \int d\rho dE_l\ [M_{\Lambda_c}\sqrt{\rho^2-1}\cos\bar{\theta}\ F_1(E', E_l) + E_l\ F_2(E', E_l)]\ . \end{aligned} \tag{23}$$

It is straightforward to show, from Eq. (22),

$$N_\pm = C_0(G_0 \pm \frac{1}{2}G_1\ \mathbf{P}\cdot\hat{e})\ , \tag{24}$$

C_0 being a constant depending on the production rate of Λ_b baryons. Therefore, we have

$$\alpha(\hat{e}) = \frac{G_1}{2G_0}\ \mathbf{P}\cdot\hat{e} = -0.31\mathbf{P}\cdot\hat{e}\ , \tag{25}$$

where the numerical results of G_0 and G_1 from Eq. (23) have been inserted. It implies that the lepton tends to be emitted in the direction opposite to the transverse polarization of the Λ_b baryon and that the magnitude of the asymmetry parameter can not exceed 0.31.

We have tested the sensitivity of our predictions to the variation of the heavy baryon wave functions. The asymmetry parameters $\alpha(\hat{e})$ change only few percents for various choices of the baryon wave functions. This observation is expected, since nonperturbative effects from the wave functions cancel in the ratio of $(N_+ - N_-)/(N_+ + N_-)$. For a similar reason, this ratio does not depend on the normalizations of the heavy baryon wave functions. Therefore, Eq. (25) can be regarded as being almost model-independent, and verified experimentally. Other potential corrections to the asymmetry parameter are under control and can be included systematically. For example, higher-order corrections to the hard amplitudes are suppressed by $\alpha_s/\pi < 0.3$, which give 30% uncertainty at most. Higher-twist contributions are of order $\bar{\Lambda}/M_{\Lambda_b} \sim 0.1$ with the mass difference $\bar{\Lambda} = M_{\Lambda_b} - m_b$, m_b being the b quark mass.

The $\bar{\Lambda}_b$ baryon, produced together with the Λ_b baryon in proton-anti-proton collisions, possesses a polarization opposite to that of the Λ_b baryon, while the lepton from the $\bar{\Lambda}_b \to \bar{\Lambda}_c \bar{l}\nu$ decay tends to be produced along the transverse polarization of the $\bar{\Lambda}_b$ baryon. As a result, for colliding experiments at the Tevatron, it is possible to combine events of the Λ_b and $\bar{\Lambda}_b$ baryons to double the statistics, and measure the lepton asymmetry parameter irrespective of whether the leptons are from Λ_b or $\bar{\Lambda}_b$ baryon decays. With about 200 Λ_b baryon events available currently, the asymmetry parameter could be measured to 10% accuracy corresponding to a 30% polarization. For Run II of the Tevatron, it is likely to collect 20 times more events so that a polarization of a few percent could be determined. In addition, the Λ_b baryon polarization can be measured through meson asymmetry parameter in nonleptonic decays. The application of the PQCD formalism to this case is under investigation.

III. CONCLUSION

In this paper we have studied the polarized Λ_b baryon semileptonic decay $\Lambda_b \to \Lambda_c l \bar{\nu}$ using PQCD factorization theorem. We have proposed a boost invariant quantity $\alpha(\hat{e})$ related to the transverse polarization $\mathbf{P} \cdot \hat{e}$ of a Λ_b baryon, which can be measured with available data from current and future experiments. This quantity is defined as the number asymmetry of the leptons which are produced with positive and negative components of momenta along $\hat{e}$. With respect to the production plane of Λ_b baryons at the Tevatron, it is the asymmetry of the lepton numbers above and below this plane. Using the transition form factors obtained in the PQCD approach, we have predicted $\alpha(\hat{e}) = -0.31\mathbf{P} \cdot \hat{e}$. It indicates that the lepton tends to be emitted in the direction opposite to the Λ_b baryon transverse polarization and that the magnitude of the asymmetry parameter can not exceed 0.31. Our result is useful for extracting the transverse polarization of a Λ_b baryon from its exclusive semileptonic decay.

This work was supported by the National Science Council of the Republic of China under Grant Nos. NSC-88-2112-M-001-041 and NSC-89-2112-M-006-004.

[1] H.H. Shih, S.C. Lee, and H-n. Li, Phys. Rev. D **59**, 094014 (1999).
[2] H.H. Shih, S.C. Lee, and H-n. Li, hep-ph/9906370, to appear in Phys. Rev. D.
[3] The ALEPH Collaboration, D. Buskulic *et al.*, Phys. Lett. B **365**, 437 (1996).
[4] The OPAL Collaboration, G. Abbiendi *et al.*, Phys. Lett. B **444**, 539 (1998).
[5] G. Bonvicini and L. Randall, Phys. Rev Lett. **73**, 392 (1994).
[6] C. Diaconu, M. Talby, J.G. Körner, and D. Pirjol, Phys. Rev. D **53**, 6186 (1996).
[7] N. Isgur and M.B. Wise, Phys. Lett. B **232**, 113 (1989); **237**, 527 (1990).
[8] Particle Data Group, C. Caso *et al.*, Euro. Phys. J. C **3**, 1 (1998).
[9] X.H. Guo and P. Kroll, Z. Phys. C **59**, 567 (1993); B. Konig, J.G. Korner, M. Kramer, and P. Kroll, Phys. Rev. D **56**, 4282 (1997).
[10] B. König, J.G. Körner, M. Krämer, and P. Kroll, Phys. Rev. D **56**, 4282 (1997).
[11] M.A. Ivanov, J.G. Körner, V.E. Lyubovitskij, and A.G. Rusetsky, Report No. hep-ph/9910342 (and references therein).
[12] M.A. Ivanov, J.G. Körner, V.E. Lyubovitskij, and A.G. Rusetsky, Phys. Rev. D **59**, 074016 (1999).
[13] H.G. Dosch, E. Ferreira, M. Nielsen, and R. Rosenfeld, Phys. Lett. B **431**, 173 (1998).
[14] R.S.M. de Carvalho, F.S. Navarra, M. Nielsen, E. Ferreira, and H.G. Dosch, Phys. Rev. D **60**, 034009 (1999).
[15] Y.B. Dai, C.S. Huang, M.Q. Huang and C. Liu, Phys. Lett. B **387**, 379 (1996).
[16] S.C. Lee, Phys. Lett. B **189**, 461 (1987).
[17] H. Georgi, B. Grinstein and M.B. Wise, Phys. Lett. B **252**, 456 (1990).
[18] A.F. Falk and M. Neubert, Phys. Rev. D **47**, 2982 (1993).

C	0.2	0.3	0.4	0.5
$\rho = 1.2$	18%	49%	66%	78%
$\rho = 1.3$	22%	54%	71%	79%
$\rho = 1.4$	25%	58%	74%	82%

TABLE I. Percentages of the full contributions to the form factor f_1, that arise from the region with $\alpha_s/\pi < C$.